

CA: d
cont
aminopropyl)methylamine, and the LYSOTRACKER[®] probes which report intralysosomal pH as well as the dynamic distribution of lysosomes (Molecular Probes, Inc.)

Please replace the text at page 101 lines 10-14 with the following:

Mitochondrial labeling

C10
In one embodiment, membrane permeant mitochondrial-specific luminescent reagents (Molecular Probes, Inc.) are used to label the mitochondria of living and fixed cells. These reagents include rhodamine 123, tetramethyl rosamine, JC-1, and the MITOTRACKER[®] reactive dyes.

In the claims:

Please cancel claims 1 and 18

Please add the following new claims:

- Sub-101
R-126
211
19. (NEW)
- C11
- a) providing an array of locations comprising cells, wherein the cells possess at least a first luminescently labeled reporter molecule that reports on cell location, and at least a second luminescently labeled reporter molecule that reports on neurite outgrowth;
- b) obtaining a nuclear image from the at least first luminescently labeled reporter molecule and a neurite image from the at least second luminescently-labeled reporter molecule;
- c) automatically identifying cell bodies from the nuclear image;
- d) automatically identifying neurites extending from the cell bodies from the neurite image; and
- e) automatically determining one or more neurite features selected from the group consisting of:
- i) Total neurite length from all cells;
 - ii) Total number of neurite branches from all cells;
 - iii) Number of neurites per cell;
 - iv) Number of neurites per positive neuron;
 - v) Neurite length from each cell;
 - vi) Neurite length per positive neuron;
 - vii) Neurite length per neurite;

- viii) Number of cells that are positive for neurite outgrowth;
- ix) Percentage of cells positive for neurite outgrowth;
- x) Number of branches per neuron; and
- xi) Number of branches per neurite.

²²
20. (NEW) The method of claim 19 wherein identifying cell bodies comprises the steps of:

- A) generating a kernel image from the nuclear image;
- B) performing conditional dilations of the kernel image to identify the cell body.

²³
21. (NEW) The method of claim 20, wherein identifying neurites extending from cell bodies comprises the steps of:

- I) generating a reservoir image from the neurite-image; and
- II) identifying positive pixels in the reservoir image that are not present in the cell bodies, wherein such positive pixels belong to neurites extending from cell bodies.

²⁴
22. (NEW) The method of claim 21, further comprising

- (a) performing one conditional dilation of the kernel image to acquire a dilation image;
- (b) determining a set of nodes from the dilation image;
- (c) linking together connected nodes; and
- (d) repeating steps (a)-(c) until an entire neurite length has been traced.

²⁵
23. (NEW) The method of claim 22, wherein steps (a) through (d) are carried out at multiple time points.

²⁶
24. (NEW) The method of claim 19 further comprising contacting the neurons with a test compound, and determining an effect of the test compound on neurite outgrowth from the cell bodies.

²⁷
25. (NEW) The method of claim 24, further comprising contacting the neurons with a neurotoxin either before, after, or simultaneously with the test compound.

²⁸
26. (NEW) The method of claim 24, further comprising contacting the cells with a control compound known to stimulate neurite outgrowth, and determining whether the test compound inhibits the control compound from inducing neurite outgrowth from the cell bodies.

²⁹
27. (NEW) The method of claim 19, wherein steps b) through e) are carried out at multiple time points.

³⁰
28. (NEW) The method of claim 19 wherein the first luminescently labeled reporter molecule comprises a DNA binding compound.